

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

20040601 029

FINAL REPORT

Grant #: N00014-00-1-0314

PRINCIPAL INVESTIGATOR: Tania Zenteno-Savín, Ph.D.

GRANT TITLE: Coping with repeated ischemia and reperfusion:
Physiology of Free Radicals in Diving Birds and Mammals

AWARD PERIOD: 20 February 2000 - 19 February 2003

OBJECTIVE: The specific mechanisms involved in protecting marine bird and mammal tissues from generation of free radicals and their deleterious effects are not known and are the subject of this research. The short-term objectives of this study are to analyze the rate of production of superoxide radical as an index of synthesis of free radicals, in tissues from marine birds and mammals; to examine the degree of lipid peroxidation as an index of free radical-induced damage, in tissues from marine birds and mammals; and to do a selective analysis of the antioxidant defenses which may endow marine birds and mammals with the tolerance to repeated cycles of ischemia/reperfusion associated to diving.

APPROACH: Using colorimetric techniques, we are evaluating the rate of production of superoxide radicals, lipid peroxidation, and the activities of the main antioxidant enzymes in extracts of tissues from ringed seals, *Phoca hispida*, and emperor penguins *Aptenodytes forsteri*. These determinations are made under basal conditions and under simulated oxidative stress.

ACCOMPLISHMENTS: Baseline production of superoxide radical ($O_2^{\cdot -}$) and lipid peroxidation (TBARS) were obtained from 8 emperor penguins, 3 northern elephant seals and 3 California sea lions; data is available for swimming (S) and non-swimming (NS) muscle biopsies from these species. These results were obtained in collaboration with Dr. Paul Ponganis at the University of California San Diego-Scripps Institution of Oceanography. We have analyzed the production of $O_2^{\cdot -}$, TBARS and total antioxidant capacity (AOX) in samples of heart, kidney and muscle from ringed seals, that have been obtained with the cooperation of alaskan subsistence hunters. Samples from 17 seals were obtained and analyzed, and compared with results obtained from 13 pigs as a model of terrestrial mammal. Oxidative stress was simulated in these samples by incubating with xanthine/xanthine oxidase (X/XO). Activity of the antioxidant enzymes catalase, superoxide

dismutase and glutathione S-transferase were analyzed in tissues samples from ringed seals. These results are product of collabrations with Dr Robert Elsner, University of Alaska Fairbanks. Additionally, in collaboration with Dr. Peter Johnson, Ohio University, we are studying the presence of hipoxia-inducible factor (HIF) in ringed seal tissues.

$O_2^{\cdot -}$ production in ringed seal and elephant seal was higher in S than NS muscle ($P < 0.1$). $O_2^{\cdot -}$ production in California sea lion and penguin samples was higher in NS than in S muscle ($P < 0.1$). $O_2^{\cdot -}$ production was higher in all marine species than in domestic pig ($P < 0.1$). TBARS were higher in NS than S muscle in California sea lion ($P < 0.1$). Emperor penguin S muscle had higher TBARS than NS muscle ($P < 0.1$). While seal, sea lion and penguin muscle showed much higher $O_2^{\cdot -}$ production than domestic pig muscle, tissue damage, as evidenced by TBARS levels, was not proportionally increased in the marine bird and mammals. Antioxidant capacity was analyzed in ringed seal and domestic pig muscle samples, and was found to be higher in ringed seal ($P < 0.1$).

CONCLUSIONS: In reponse to cyclic episodes of ischemia/reperfusion associated with their frequent dives, one might expect that seal, sea lion and penguin tissues have biochemical and/or physiological adaptations for tolerance of, or protection from, the potential oxidative insult. In this study we found higher production of $O_2^{\cdot -}$ in muscle from ringed seal, elephant seal, California sea lion and emperor penguin than in muscle from the domestic pig. Despite the higher $O_2^{\cdot -}$ production found in penguin, seals and sea lion, tissue damage was not proportionally increased in the marine bird and mammals, as evidenced by TBARS levels. The differences in AOX between pig and ringed seal muscle, provide a suitable explanation for this protection against $O_2^{\cdot -}$. Differences between S and NS may be due to fiber type distribution. Alternatively, differences among species may be related to diving capacities. Overall, the results from this study suggest that tolerance of dive-associated ischemia/reperfusion in marine organisms may depend on enhanced intermediate scavenging of reactive oxygen species. Oxidative stress was simulated in these samples by incubating with xanthine/xanthine oxidase (X/XO). Treatment with X/XO had no effect on TBARS in ringed seal kidney or heart. These results show that ringed seal muscle, heart and kidney can be induced *in vitro* to generate reactive oxygen species, and suggest that the living seal's protective defenses may depend on $O_2^{\cdot -}$ production, similar to the protective effect of experimental preconditioning, or on enhanced intermediate scavenging, as evidenced by the larger AOX found in ringed seal tissues compared to pig tissues. Additionally, activity of the antioxidant enzymes catalase,

superoxide dismutase and glutathione S-transferase were analyzed in tissues samples from ringed seals. These results are in accordance to those obtained by other researchers (e.g., Wilhelm-Filho et al., 2002), and suggest that marine mammals have an enhanced antioxidant capacity that allows them tolerance of the potential oxidative stress associated to repeated cycles of ischemia-reperfusion.

SIGNIFICANCE: This project seeks to understand the metabolic adaptations and physiologic mechanisms that allow marine birds and mammals to cope with the hypoxia, ischemia/reperfusion and oxidative stress associated to breath-hold diving, which would help us understand how this factors affect human divers and how to deal with specific pathologies associated with exposure to hypoxia, ischemia/reperfusion and oxidative stress. For example, oxygen toxicity, an important problem for human divers, is at the cellular level an effect of the free radicals; therefore, marine birds and mammals can be thought of as models for the study of such condition.

PUBLICATIONS AND ABSTRACTS (for total period of grant):

Zenteno-Savín, T. & R. Elsner (2000) "Differential oxidative stress in ringed seals", Free Rad. Biol. & Med. 29 (Suppl 1): S39.

7th Annual Meeting of The Oxygen Society- The Oxygen Society. San Diego, CA, U.S.A. Noviembre 2000. Zenteno-Savín, T. y Robert Elsner. "Differential oxidative stress in ringed seal tissues". XXVI Reunión Internacional para el Estudio de los Mamíferos Marinos en México- SOMEMMA (Sociedad Mexicana de Mastozoología Marina, A.C.). Ensenada, Baja California, México. Mayo, 2001. Clayton Hernández, E., T. Zenteno-Savín, y R. Elsner: "Producción y efecto de los radicales libres en tejidos de foca anillada". Workshop on Comparative Aspects of Oxidative Stress in Biological Systems. La Paz, Baja California Sur. Octubre 16-19, 2001. T. Zenteno-Savín, E. Clayton Hernández, y R. Elsner. "Diving seals, are they a model for coping with oxidative stress?".

14th Biennial Conference on the Biology of Marine Mammals. Vancouver, B.C., Canadá. Noviembre 28- Diciembre 3, 2001. T. Zenteno-Savín, E. Clayton Hernández y R. Elsner. "Comparative oxidative stress: Ringed seals and pigs". M. Hermes-Lima y T. Zenteno-Savín. 2002. Animal response to drastic changes in oxygen availability and physiological stress. Review Article. Comparative Biochemistry and Physiology. 133: 537-556.

Zenteno-Savín, T., E. Clayton-Hernández & R. Elsner (2002) "Diving seals: Are they a model for coping with oxidative stress?", Comp Biochem Physiol, 133: 527-536.

Zenteno-Savín, T. (2002) "Oxidative stress in marine organisms: A review". In: Oxidative Stress at Molecular, Cellular and Organ Levels. P. Johnson and A.A. Boldyrev, Eds. Research Signpost, Kerala, India. Pp. 67-76. Book chapter.

2002 Alaskan Summer Neuroscience Conference. Fairbanks, AK, USA. July 14-15, 2002. T. Zenteno-Savín. "Antioxidant protection in marine birds and mammals"

2002 Alaskan Summer Neuroscience Conference. Fairbanks, AK, USA. July 14-15, 2002. M. Hermes-Lima, T. Zenteno-Savín, G. Mourão. "Adaptations against oxidative stress in hypoxia-tolerant animals".

XLV Congreso Nacional de Ciencias Fisiológicas. Colima, Mexico, 8-12 September, 2002. T. Zenteno-Savín & R. Elsner. "Potencial estrés oxidativo en focas anilladas".

The Power of Comparative Physiology. San Diego, CA. August 24-28, 2002. T. Zenteno-Savín, R. Elsner & P. Ponganis. "Antioxidant protection in marine birds and mammals".

Johnson, P., T. Zenteno-Savín & R. Elsner (2003) "Hipoxia-inducible factor in ringed seal tissues". Free Rad. Biol. & Med. 35 (Suppl 1): S77.

XLVI Congreso Nacional de Ciencias Fisiológicas. Aguascalientes, Mexico. 3-6 August, 2003. T. Zenteno-Savín. "Estrés oxidativo en organismos marinos".

XXXVIII Congresso da Sociedade Brasileira de Fisiologia -SBFis y XXI Congresso da Associação Latino Americana de Ciências Fisiológicas -ALACF. Riberão Preto, São Paulo, Brasil. 1-4 Septiembre, 2003. T. Zenteno-Savín. "Potential oxidative stress in diving vertebrates" invited speaker to the Symposium "Metabolic depression and life in the limit of oxygenation".

10th Annual Meeting of The Oxygen Society- Society for Free Radical Biology and Medicine. Seattle, WA, U.S.A. Noviembre 20-24, 2003. P. Johnson, Zenteno-Savín, T. y R. Elsner. "Hipoxia-inducible factor in ringed seal tissues".

Johnson, P., Elsner, R., Zenteno-Savín, T. 2004. Sequence analysis of hypoxia-inducible factor -1 * from ringed seal. Free Radical Biology and Medicine, 36(Suppl1): S54.

XXIX Reunión Internacional para el Estudio de los Mamíferos Marinos. La Paz, BCS, Mayo 2-5, 2004. Vázquez-Medina, J.P., Zenteno-Savín, T., Elsner, R. "Actividad enzimática antioxidante en tejidos de foca anillada (*Phoca hispida*) y cerdo común (*Sus scrofa*)".

12th Biennial Meeting of the Society for Free Radical Research International. Buenos Aires, Argentina, Mayo 5-9, 2004. Johnson, P., Elsner, R., Zenteno-Savín, T. "Sequence analysis of hypoxia-inducible factor -1 α from ringed seal".

Johnson, P., T. Zenteno-Savín y R. Elsner. HIF in ringed seal tissues. Free Radicals in Biology and Medicine. In press.